




Rabbit Anti-Goat IgG Fab

Cat. No.	Format	Size
6022-01	Purified (UNLB)	0.5 mg
6022-08	Biotin (BIOT)	0.5 mg



ELISA plate was coated with purified goat IgG Fab and IgG Fc. Immunoglobulins were detected with serially diluted Rabbit Anti-Goat IgG Fab-UNLB (SB Cat. No. 6022-01) followed by Goat Anti-Rabbit IgG(H+L), Mouse/Human ads-HRP (SB Cat. No. 4050-05).

Description

Specificity	Reacts with the Fab region of goat IgG
Source	Pooled antisera from rabbits hyperimmunized with goat IgG
Cross Adsorption	Goat IgG Fc; may react with immunoglobulins from other species and the light chains of other goat immunoglobulins
Purification	Affinity chromatography on goat IgG Fab covalently linked to agarose

Applications

Quality tested applications include –
ELISA

Working Dilutions

ELISA	Purified (UNLB) antibody	≤ 1 µg/mL
	BIOT conjugate	1:5,000 – 1:20,000
Other Applications	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

TB6022
09-Dec-19