




Rabbit Anti-Goat IgG(H+L)

Cat. No.	Format	Size
6160-01	Purified (UNLB)	1.0 mg
6160-02	Fluorescein (FITC)	1.0 mg
6160-03	Rhodamine (TRITC)	1.0 mg
6160-04	Alkaline Phosphatase (AP)	1.0 mL
6160-05	Horseradish Peroxidase (HRP)	1.0 mL
6160-08	Biotin (BIOT)	1.0 mg



Paraffin embedded rat kidney section post 30% blood volume loss and treatment with ketoprofen was stained with anti-IL-1 followed by Rabbit Anti-Goat IgG(H+L)-BIOT (SB Cat. No. 6160-08), HRP conjugated streptavidin, DAB, and hematoxylin.

Image from Guedes FS Jr, Cruz DS, Rodrigues MM, Silva LM, Amorim RL, Viana PT, et al. Renal histology and immunohistochemistry after acute hemorrhage in rats under sevoflurane and ketoprofen effect. Acta Cir Bras. 2012;27:37-42. Figure 3

Description

Specificity	Reacts with the heavy and light chains of goat IgG
Source	Pooled antisera from rabbits hyperimmunized with goat IgG
Cross Adsorption	None; may react with immunoglobulins from other species and the light chains of other goat immunoglobulins
Purification	Affinity chromatography on goat IgG covalently linked to agarose

Applications

Quality tested applications include –

ELISA¹⁻⁷
FLISA

Other referenced applications include –

IHC-FS⁸⁻¹⁰
IHC-PS^{11,12}
ICC^{13,14}
WB¹⁴⁻²¹

Working Dilutions

ELISA	AP conjugate	1:2,000 – 1:4,000
	HRP conjugate	1:4,000 – 1:8,000
	BIOT conjugate	1:5,000 – 1:20,000
FLISA	FITC conjugate	1:200 – 1:400
	TRITC conjugate	1:100 – 1:400
Other Applications	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added. Store at 2-8°C.
- The fluorescein (FITC) and rhodamine (TRITC) conjugates are supplied as 1.0 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50 mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN₃. Store at 2-8°C.
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

- Costa-Pinto FA, Basso AS, Britto LR, Malucelli BE, Russo M. Avoidance behavior and neural correlates of allergen exposure in a murine model of asthma. *Brain Behav Immun.* 2005;19:52-60. (ELISA)
- Costa-Pinto FA, Basso AS, Russo M. Role of mast cell degranulation in the neural correlates of the immediate allergic reaction in a murine model of asthma. *Brain Behav Immun.* 2007;21:783-90. (ELISA)
- Shi W, Li YH, Liu F, Yang JY, Zhou DH, Chen YQ, et al. Flagellin enhances saliva IgA response and protection of anti-caries DNA vaccine. *J Dent Res.* 2012;91:249-54. (ELISA)
- Will TA, Polcari AJ, Garcia JG, Ouwenga MK, Voelzke BB, Greisler HP, et al. Paclitaxel inhibits ureteral smooth muscle cell proliferation and collagen production in the absence of cell toxicity. *J Urol.* 2011;185:335-40. (ELISA)
- Jain S, Singh SR, Horn DW, Davis VA, Ram MK, Pillai S. Development of an antibody functionalized carbon nanotube biosensor for foodborne bacterial pathogens. *J Biosens Bioelectron.* 2012;S11:002. (ELISA)
- Lima FA, Miyaji EN, Quintilio W, Raw I, Ho PL, Oliveira ML. Pneumococcal Surface Protein A does not affect the immune responses to a combined diphtheria tetanus and pertussis vaccine in mice. *Vaccine.* 2013;31:2465-70. (ELISA)
- Faustino L, Fonseca DM, Florsheim EB, Ressende RR, Lepique AP, Faquim-Mauro E, et al. Tumor necrosis factor-related apoptosis-inducing ligand mediates the resolution of allergic airway inflammation induced by chronic allergen inhalation. *Mucosal Immunol.* 2014;7:1199-208. (ELISA)
- Li Y, Qin X, Kuang F, Wu R, Duan X, Ju G, et al. Alterations of Fc gamma receptor I and Toll-like receptor 4 mediate the antiinflammatory actions of microglia and astrocytes after adrenaline-induced blood-brain barrier opening in rats. *J Neurosci Res.* 2008;86:3556-65. (IHC-FS)
- Xia RH, Yosef N, Ubogu EE. Selective expression and cellular localization of pro-inflammatory chemokine ligand/receptor pairs in the sciatic nerves of a severe murine experimental autoimmune neuritis model of Guillain-Barré syndrome. *Neuropathol Appl Neurobiol.* 2010;36:388-98. (IHC-FS)
- Jansen ID, Hollander AP, Buttle DJ, Everts V. Type II and VI collagen in nasal and articular cartilage and the effect of IL-1 α on the distribution of these collagens. *J Mol Hist.* 2010;41:9-17. (IHC-FS)
- Yoon A, Kim J, Lee Y, Kim H, Yoo J, Sohn J, et al. Markedly enhanced cytolysis by E1B-19kD-deleted oncolytic adenovirus in combination with cisplatin. *Hum Gene Ther.* 2006;17:379-90. (IHC-PS)
- Guedes FS Jr, Cruz DS, Rodrigues MM, Silva LM, Amorim RL, Vianna PT, et al. Renal histology and immunohistochemistry after acute hemorrhage in rats under sevoflurane and ketoprofen effect. *Acta Cir Bras.* 2012;27:37-42. (IHC-PS)
- Cambridge CD, Singh SR, Waffo AB, Fairley SJ, Dennis VA. Formulation, characterization, and expression of a recombinant MOMP Chlamydia trachomatis DNA vaccine encapsulated in chitosan nanoparticles. *Int J Nanomedicine.* 2013;8:1759-71. (ICC)
- Duan Y, Huang S, Yang J, Niu P, Gong Z, Liu X, et al. HspA1A facilitates DNA repair in human bronchial epithelial cells exposed to Benzo[a]pyrene and interacts with casein kinase 2. *Cell Stress Chaperones.* 2014;19:271-9. (ICC, WB)
- Webb CF, Yamashita Y, Ayers N, Evetts S, Paulin Y, Conley ME, et al. The transcription factor Bright associates with Bruton's tyrosine kinase, the defective protein in immunodeficiency disease. *J Immunol.* 2000;165:6956-65. (WB)
- Zhang C, Yang F, Zhang X, Wang S, Li M, Lin L, et al. Grateloupia longifolia polysaccharide inhibits angiogenesis by downregulating tissue factor expression in HMEC-1 endothelial cells. *Br J Pharmacol.* 2006;148:741-51. (WB)
- Røsjø H, Husberg C, Dahl MB, Stridsberg M, Sjaastad I, Finsen AV, et al. Chromogranin B in heart failure: a putative cardiac biomarker expressed in the failing myocardium. *Circ Heart Fail.* 2010;3:503-11. (WB)
- Lunde IG, Kvaløy H, Austbø B, Christensen G, Carlson CR. Angiotensin II and norepinephrine activate specific calcineurin-dependent NFAT transcription factor isoforms in cardiomyocytes. *J Appl Physiol.* 2011;111:1278-89. (WB)
- Shibasaki M, Watanabe K, Takeda K, Itoh T, Tsuyuki T, Narita M, et al. Effect of chronic ethanol treatment on μ -opioid receptor function, interacting proteins and morphine-induced place preference. *Psychopharmacology.* 2013;228:207-15. (WB)
- Lou Y, Wu C, Wu M, Xie C, Ren L. The changes of neutrophil gelatinase-associated lipocalin in plasma and its expression in adipose tissue in pregnant women with gestational diabetes. *Diabetes Res Clin Pract.* 2014;104:136-42. (WB)
- Lee WJ, Ahn HM, Roh H, Na Y, Choi I, Lee JH, et al. Decorin-expressing adenovirus decreases collagen synthesis and upregulates MMP expression in keloid fibroblasts and keloid spheroids. *Exp Dermatol.* 2015 Apr 11. doi: 10.1111/exd.12719. [Epub ahead of print]. (WB)

TB6160
28-Oct-19