




Goat Anti-Human Ig

Cat. No.	Format	Size
2010-01	Purified (UNLB)	2.0 mg
2010-02	Fluorescein (FITC)	1.0 mg
2010-03	Rhodamine (TRITC)	1.0 mg
2010-04	Alkaline Phosphatase (AP)	1.0 mL
2010-05	Horseradish Peroxidase (HRP)	1.0 mL
2010-06	β -galactosidase (BGAL)	1.0 mL
2010-07	Texas Red [®] (TXRD)	1.0 mg
2010-08	Biotin (BIOT)	1.0 mg
2010-09	R-phycoerythrin (PE)	0.5 mg
2010-30	Alexa Fluor [®] 488 (AF488)	1.0 mg
2010-31	Alexa Fluor [®] 647 (AF647)	1.0 mg
2010-32	Alexa Fluor [®] 555 (AF555)	1.0 mg



Human peripheral blood lymphocytes were stained with Goat Anti-Human Ig-PE (SB Cat. No. 2010-09) and Mouse Anti-Human CD19-FITC (SB Cat. No. 9340-02).

Description

Specificity	Reacts with the heavy and light chains of human IgG, IgM, and IgA
Source	Pooled antisera from goats hyperimmunized with human IgG, IgM, and IgA
Cross Adsorption	None; may react with immunoglobulins from other species
Purification	Affinity chromatography on pooled human immunoglobulins covalently linked to agarose

Applications

Quality tested applications include –

ELISA¹⁻⁸
 FLISA
 FC^{15,16,18}

Other referenced applications include –

ICC⁹
 WB¹⁰⁻¹³
 Multiplex¹⁴
 Stimulation¹⁵⁻¹⁷

Working Dilutions

ELISA	AP conjugate	1:2,000 – 1:4,000
	HRP conjugate	1:4,000 – 1:8,000
	BGAL conjugate	1:500
	BIOT conjugate	1:5,000 – 1:20,000
FLISA	FITC, TRITC, TXRD, AF488, and AF555 conjugates	1:100 – 1:400
	PE and AF647 conjugates	$\leq 1 \mu\text{g/mL}$
Flow Cytometry	FITC, BIOT, and AF488 conjugates	$\leq 1 \mu\text{g}/10^6 \text{ cells}$
	PE and AF647 conjugates	$\leq 0.1 \mu\text{g}/10^6 \text{ cells}$
	For flow cytometry, the suggested use of these reagents is in a final volume of 100 μL	
Other Applications	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 2.0 mg purified immunoglobulin in 2.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC), rhodamine (TRITC), Texas Red® (TXRD), Alexa Fluor® 488 (AF488), Alexa Fluor® 555 (AF555), and Alexa Fluor® 647 (AF647) conjugates are supplied as 1.0 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50 mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The β-galactosidase (BGAL) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Kouzmitcheva GA, Petrenko VA, Smith GP. Identifying diagnostic peptides for Lyme disease through epitope discovery. *Clin Diagn Lab Immunol.* 2001;8:150-60. (ELISA)
2. Pancook JD, Beuerlein G, Pecht G, Tang Y, Nie Y, Wu H, et al. In vitro affinity maturation of human IgM antibodies reactive with tumor-associated antigens. *Hybrid Hybridomics.* 2001;20:383-96. (ELISA)
3. Weitkamp J, Kallewaard N, Kusahara K, Feigelstock D, Feng N, Greenberg HB, et al. Generation of recombinant human monoclonal antibodies to rotavirus from single antigen-specific B cells selected with fluorescent virus-like particles. *J Immunol Methods.* 2003;275:223-37. (ELISA)
4. Ohdan H, Zhou W, Tanaka Y, Irei T, Fuchimoto Y, Egawa H, et al. Evidence of immune tolerance to blood group antigens in a case of ABO-incompatible pediatric liver transplantation. *Am J Transplant.* 2007;7:2190-4. (ELISA)
5. Williams JV, Weitkamp J, Blum DL, LaFleur BJ, Crowe JE Jr. The human neonatal B cell response to respiratory syncytial virus uses a biased antibody variable gene repertoire that lacks somatic mutations. *Mol Immunol.* 2009;47:407-14. (ELISA)
6. Fan Q, Longnecker R. The Ig-like V-type domain of paired Ig-like type 2 receptor alpha is critical for herpes simplex virus type 1-mediated membrane fusion. *J Virol.* 2010;84:8664-72. (ELISA)
7. Yang G, Holl TM, Liu Y, Li Y, Lu X, Nicely NI, et al. Identification of autoantigens recognized by the 2F5 and 4E10 broadly neutralizing HIV-1 antibodies. *J Exp Med.* 2013;210:241-56. (ELISA)
8. Marti GP, Mohn KG, Cox RJ, Brokstad KA. The influence of tonsillectomy on total serum antibody levels. *Scand J Immunol.* 2014;80:377-9. (ELISA)
9. Moody MA, Yates NL, Amos JD, Drinker MS, Eudailey JA, Gurley TC, et al. HIV-1 gp120 vaccine induces affinity maturation in both new and persistent antibody clonal lineages. *J Virol.* 2012;86:7496-507. (ICC)
10. Moritz DR, Gheyselincx J, Klemenz R. Expression analysis of the soluble and membrane-associated forms of the interleukin-1 receptor-related TI protein in primary mast cells and fibroblasts. *Hybridoma.* 1998;17:107-16. (WB)
11. Davenport V, Guthrie T, Findlow J, Borrow R, Williams NA, Heyderman RS. Evidence for naturally acquired T cell-mediated mucosal immunity to *Neisseria meningitidis*. *J Immunol.* 2003;171:4263-70. (WB)
12. Botelho-Nevers E, Gouriet F, Lepidi H, Couvret A, Amphoux B, Dessi P, et al. Chronic nasal infection caused by *Klebsiella rhinoscleromatis* or *Klebsiella ozaenae*: two forgotten infectious diseases. *Int J Infect Dis.* 2007;11:423-9. (WB)
13. Parcina M, Miranda-Garcia MA, Durlanik S, Ziegler S, Over B, Georg P, et al. Pathogen-triggered activation of plasmacytoid dendritic cells induces IL-10-producing B cells in response to *Staphylococcus aureus*. *J Immunol.* 2013;190:1591-602. (WB)
14. Pochechueva T, Chinarev A, Bovin N, Fedier A, Jacob F, Heinzlmann-Schwarz V. PEGylation of microbead surfaces reduces unspecific antibody binding in glycan-based suspension array. *J Immunol Methods.* 2014;412:42-52. (Multiplex)
15. Fruehling S, Lee SK, Herrold R, Frech B, Laux G, Kremmer E, et al. Identification of latent membrane protein 2A (LMP2A) domains essential for the LMP2A dominant-negative effect on B-lymphocyte surface immunoglobulin signal transduction. *J Virol.* 1996;70:6216-26. (FC, Stim)
16. Fruehling S, Swart R, Dolwick KM, Kremmer E, Longnecker R. Tyrosine 112 of latent membrane protein 2A is essential for protein tyrosine kinase loading and regulation of Epstein-Barr virus latency. *J Virol.* 1998;72:7796-806. (FC, Stim)
17. Koethe S, Zander L, Köster S, Annan A, Ebenfelt A, Spencer J, et al. Pivotal advance: CD45RB glycosylation is specifically regulated during human peripheral B cell differentiation. *J Leukoc Biol.* 2011;90:5-19. (Stim)
18. Wiehr S, Bühler P, Gierschner D, Wolf P, Rolle A, Kesenheimer C, et al. Pharmacokinetics and PET imaging properties of two recombinant anti-PSMA antibody fragments in comparison to their parental antibody. *Prostate.* 2014;74:743-55. (FC)

Texas Red® is a registered trademark of Molecular Probes, Inc.

Alexa Fluor® 488, 647, and 555 are provided under an Intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.