

Certificate of Analysis

A peptide corresponding to acetyl-Histone H2A.X (Lys9). Protein A affinity purified from an animal origin–free culture supernatant 100 μg 1.0 mg/mL 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide WB: 0.01 μg/mL – 0.1 μg/mL; Multiplex: 0.2 μg/mL - 1 μg/mL. Stable for 1 Year at -20.0°C from date of receipt. U.S.A.
Protein A affinity purified from an animal origin–free culture supernatant 100 µg 1.0 mg/mL 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide WB: 0.01 µg/mL – 0.1 µg/mL; Multiplex: 0.2 µg/mL - 1 µg/mL.
Protein A affinity purified from an animal origin–free culture supernatant 100 μg 1.0 mg/mL 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide WB: 0.01 μg/mL – 0.1 μg/mL;
Protein A affinity purified from an animal origin–free culture supernatant 100 μg 1.0 mg/mL
Protein A affinity purified from an animal origin–free culture supernatant 100 μg
Protein A affinity purified from an animal origin–free culture supernatant
Protein A affinity purified from an animal origin–free
A peptide corresponding to acetyl-Histone H2A.X (Lys9).
Western Blot, Multiplex
This antibody reacts to Histone H2A.X acetylated at Lysine 9 (K9ac). No cross reactivity with non-modified Lysine 9 or other acetylated Lysines in histone H2A.
RM446
31-1337-00
Anti-Acetyl-Histone H2A.X (Lys9) Rabbit Monoclonal Antibody, Clone RM446
Rabbit Monoclonal Antibody

Therapeutic Use



Western Blot of (1) H2A recombinant protein and (2) acid extracts of HeLa cell lysate, using acetyl-Histone H2A.X (Lys9) rabbit monoclonal antibody (RM446) at 0.01 μ g/mL.



RM446 specifically reacts to Histone H2AX acetylated at Lysine 9 (K9ac). No limited cross reactivity with acetylated Lysine 5 (K5ac), Lysine 36 (K36ac) in Histone H2AX, Lysine 4 (K4ac), Lysine 7 (K7ac), Lysine 11 (K11ac), Lysine 13 (K13ac) in Histone H2AZ, or non-modified Asparagine 19 (N19), Lysine 36 (K36) in Histone H2AX.